

Oua (Uo, Nie, Uoa) Islet Bird Checklist (Between Lifou & Maré) New Caledonia

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

[You are welcome to communicate, just re-type above address into your e-mail program]

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Black-winged Petrel	<i>Pterodroma nigripennis</i>	Res bre	Br prior to 1979, 51-150 pr bred 1993-99,	1,2,
2.	Tahiti Petrel	<i>Pseudobulweria rostrata</i>	Res bre	>5 pr bred 1993-1999,	2,
3.	Wedge-tailed Shearwater	<i>Ardenna pacificus</i>	Res bre	5,500-12,000 br prs 1993-99,	2,

KEY

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

R = rare, **S** = scarce, **UC** = uncommon, **MC** = moderately common, **LC** = locally common, **C** = common, **VC** = very common, **A** = abundant, **P** = present.

Colour coding applies only to one species on a particular list. eg. An abundance comment in red for the Brown Booby refers to the Ref # in Red for that species only.

Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

REFERENCES

1. Barré, N. & Dutson, G. 2000. Liste commentée des oiseaux de Nouvelle-Calédonie. *Alauda* 3: suppl. 1-48.
2. Benoit, M.P., & Bretagnolle, V. 2002. Seabirds of the Southern Lagoon of New Caledonia; Distribution, Abundance and Threats. *Waterbirds* 25(2), 202-213.