

Radio Island Bird Checklist

Chesterfield Group, New Caledonia

19 51 37s 158 27 33e

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

[You are welcome to communicate, just re-type above address into your e-mail program]

| # | Common Name | Scientific Name | Ecol. Status | Abundance | References |
|----|------------------|----------------------------------|--------------|-----------------------------------|------------|
| 1. | Masked Booby | <i>Sula dactylatra personata</i> | Res bre | Breeding c eggs & young Nov 2007. | 1 |
| 2. | Brown Booby | <i>Sula leucogaster plotus</i> | Res bre | Breeding c eggs & young Nov 2007. | 1. |
| 3. | Red-footed Booby | <i>Sula sula</i> | Red bre | Breeding c eggs & young Nov 2007. | 1. |
| 4. | Brown Noddy | <i>Anous stolidus pileatus</i> | Res bre | P Nov 2007. | 1. |

KEY

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

R = rare, **S** = scarce, **UC** = uncommon, **MC** = moderately common, **LC** = locally common, **C** = common, **VC** = very common, **A** = abundant **P** = present.

Colour coding applies only to one species on a particular list. eg. An abundance comment in **red** for the Brown Booby refers to the Ref # in **Red** for that species only.

Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

REFERENCES

1. Emeny, Max. Personal communication July 2014.