

# Cannac Islet . Bird Checklist

## Amphlett Group, (20.5 km W of Budelum) Budi Budi Atoll. P.N.G.

9 16 03.3s 153 30 23.9e

Compiled by M.K. Tarburton, Pacific Adventist University, PNG.

[swiftlet@optusnet.com.au](mailto:swiftlet@optusnet.com.au)

[To communicate please re-type above address into your e-mail program]

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Lesser Frigatebird	<i>Fregatta ariel</i>	Res	20 obs Oct 2011,	1,
2.	Brown Booby	<i>Sula leucogaster</i>	Res	100 obs Oct 2011,	1,
3.	White-bellied Sea-Eagle	<i>Haliaeetus leucogaster</i>	Res	1 obs Oct 2011,	1,
4.	Ruddy Turnstone	<i>Arenaria interpres</i>	Sum mig	11 obs Oct 2011,	1,
5.	Black Noddy	<i>Anous minutus</i>	Res	100 Obs Oct 2011,	1,
6.	Brown Noddy	<i>Anous stolidus</i>	Res	100 obs Oct 2011,	1,
7.	Bridled Tern	<i>Onychoprion anaethetus</i>	Res	50 obs Oct 2011,	1,
8.	Crested Tern	<i>Thalasseus bergii</i>	Res	20 obs Oct 2011,	1,

### KEY

#### **Ecological status**

**End** = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

#### **Abundance**

**R** = rare, **S** = scarce, **UC** = uncommon, **MC** = moderately common, **LC** = locally common, **C** = common, **VC** = very common, **A** = abundant, **P** = present.

**Colour coding** applies only to one species on a particular list. eg. An abundance comment in **red** for the Brown Booby refers to the Ref # in **Red** for that species only.

**Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.**

### REFERENCES

1. Anonymous eBirder 11.10.2011. <http://ebird.org/ebird/view/checklist?subID=S18031554> Accessed 17.6.2015.