

Kaheu Islet. Bird Checklist

26 km W of Manus, P.N.G.

2 9 32s 146 16 20w

Compiled by M.K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Red-footed Booby	<i>Sula sula</i>	Res	1 in photo 2.1.2007,	1,
2.	Common Sandpiper	<i>Actitis hypoleucos</i>	Sum mig	12 in photo 2.1.2007,	1,
3.	Black-naped Tern	<i>Sterna s. sumatrana</i>	Res	3 in photo 2.1.2007,	1,
4.	Brown Noddy	<i>Anous stolidus pileatus</i>	Res bre	26+ in one photo 200 in another 2.1.2007,	1,
5.	Black Noddy	<i>Anous m. minutus</i>	Res	1,000+ in photos 2.1.2007, many brooding,	1,

KEY

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

R = rare, **S** = scarce, **UC** = uncommon, **MC** = moderately common, **LC** = locally common, **C** = common, **VC** = very common, **A** = abundant, **P** = present.

Colour coding applies only to one species on a particular list. eg. An abundance comment in **red** for the Brown Booby refers to the Ref # in **Red** for that species only.

Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

REFERENCES

1. Quémlangan, Blaise. 2012. Five photos on Panoramio on Google Earth at 2 9 32.36s 146 16 19.72e downloaded 11.8.2013.