

Narega Is. Bird Checklist

Bismarck Sea, PNG.

04 32 48s 149 06 44e

Compiled by M.K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

(for contact, re-type e-mail address)

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Streaked Shearwater	<i>Calonectris leucomelas</i>	Sum mig	100 obs 81 km to NW 13.1.1986.	1.
2.	Heinroth's Shearwater	<i>Puffinus heinrothi</i>	Res?	1 obs 81 km NNW 13.9.1990, 5 obs 80 km NNW 14.9.1990,	1.

KEY

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

R = rare, **S** = scarce, **UC** = uncommon, **MC** = moderately common, **LC** = locally common, **C** = common, **VC** = very common, **A** = abundant **P** = present.

Colour coding applies only to one species on a particular list. eg. An abundance comment in **red** for the Brown Booby refers to the Ref # in **Red** for that species only.

Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

REFERENCES

- Cheshire, N. 2010. Procellariiformes observed around Papua New Guinea including the Bismarck Archipelago from 1985 to 2007. South Australian Ornithologist 36, 9-24.