

Niakuni Islet Bird Checklist

Admiralty Group, PNG

2 27 s 147 e

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

swiflet@optusnet.com.au

[To communicate please re-type above address into your e-mail program]

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Coconut Lorikeet	<i>Trichoglossus haematodus</i>	Res	Known to be there by Karol Kisokau,	1,

KEY

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

R = rare, **S** = scarce, **UC** = uncommon, **MC** = moderately common, **LC** = locally common, **C** = common, **VC** = very common, **A** = abundant **P** = present.

Colour coding applies only to one species on a particular list. eg. An abundance comment in **red** for the Brown Booby refers to the Ref # in **Red** for that species only.

Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

REFERENCES

1. LeCroy, M., Peckover, W.S., & Kisokau, K. 1992. A population of Rainbow Lorikeets *Trichoglossus haematodus flavicens* roosting and nesting on the ground. *Emu* **92**, 187-190.