

# Poy-Yai (Parta-uw) Islet Bird Checklist

## 500m off Sth coast Manus. P.N.G.

2 11 25s 147 07 27.6e

Compiled by M.K. Tarburton, Pacific Adventist University, PNG.

[swiftlet@optusnet.com.au](mailto:swiftlet@optusnet.com.au)

[To communicate please re-type above address into your e-mail program]

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Eastern Reef Egret	<i>Egretta sacra</i>	Res	White & greys P. July	1,
2.	Coconut Lorikeet	<i>Trichoglossus haematodus</i>	Res bre	2 prs nesting in the ground July 1988,	1,
3.	Beach Kingfisher	<i>Todirhamphus saurophaga</i>	Res	1 obs Jul 1988,	1,
4.	Singing Starling	<i>Aplornis cantoroides</i>	Res	Prs & an imm obs Jul 1988.	1,
5.	Yellow-bellied Sunbird	<i>Nectarinia jugularis</i>	Res	A few seen Jul 1988,	1,

### KEY

#### Ecological status

**End** = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

#### Abundance

**R** = rare, **S** = scarce, **UC** = uncommon, **MC** = moderately common, **LC** = locally common, **C** = common, **VC** = very common, **A** = abundant, **P** = present.

**Colour coding** applies only to one species on a particular list. eg. An abundance comment in **red** for the Brown Booby refers to the Ref # in **Red** for that species only.

**Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.**

### REFERENCES

1. LeCroy, M., Peckover, W.S., & Kisokau, K. 1992. A population of Rainbow Lorikeets *Trichoglossus haematodus flavicens* roosting and nesting on the ground. *Emu* **92**, 187-190.