

# Baruku Island Systematic Bird Check List

## Solomon Islands

-9.0125897,158.6233521

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

[swiftlet@optusnet.com.au](mailto:swiftlet@optusnet.com.au)

[To communicate: please re-type e-mail address]

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	South Polar Skua	<i>Catharacta maccormicki</i>	mig	1 obs 12 km to west 18.2.1990,	1,
2.	Brown Noddy	<i>Anous stolidus</i>	Res	30 obs to west 18.2.1990,	1,
3.	Black Noddy	<i>Anous minutus</i>	Res	4 obs to west 18.2.1990,	1,
4.	Sooty Tern	<i>Onychoprion fuscata</i>	Res	50 obs to west 18.2.1990,	1,
5.	Bridled Tern	<i>Onychoprion anaethetus</i>	Res	10 obs to west 18.2.1990,	1,
6.	Black-naped Tern	<i>Stern sumatrana</i>	Res	50 obs to west 18.2.1990,	1,

### KEY

#### Ecological status

**End** = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

#### Abundance

**R** = rare, **S** = scarce, **UC** = uncommon, **MC** = moderately common, **LC** = locally common, **C** = common, **VC** = very common, **A** = abundant **P** = present.

**Colour coding** applies only to one species on a particular list. eg. An abundance comment in **red** for the Brown Booby refers to the Ref # in **Red** for that species only.

**Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.**

### REFERENCES

1. Scofield, P. 1990. <http://ebird.org/ebird/view/checklist?subID=S9167966> accessed 3.4.2014,