

# Nepani (Nupani) Island Bird Checklist

## Swallow Group, Santa Cruz Isles, Solomon Islands

10 02 48s 165 43 14e

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

[swiftlet@optusnet.com.au](mailto:swiftlet@optusnet.com.au)

[To communicate: please re-type e-mail address]

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Wilson's Storm Petrel	<i>Oceanites oceanicus</i>	Vag	1 coll,	8,
2.	Red-footed Booby	<i>Sula sula</i>	Res bre	said by locals on Pileni to be the source of their pet birds,	9,
3.	Eastern Reef Egret	<i>Egretta sacra</i>	Res	P, 1 spec AMNH.	1,5,
4.	Striated Heron	<i>Butorides striatus</i>	Res bre	P,	1,
5.	Pacific Golden Plover	<i>Pluvialis fulva</i>	Sum Mig	1 spec Oct	3,
6.	Pacific Imperial Pigeon	<i>Ducula pacifica pacifica</i>	Res bre	P, Coll.	1,7,
7.	Pacific Emerald Dove	<i>Chalcophaps longirostris sandwichensis</i>	Res bre	P, 1M Coll.	1,7,
8.	Coconut Lorikeet	<i>Trichoglossus haematodus</i>	Res bre	Reported by locals,	9,
9.	Long-tailed Koel	<i>Eudynamys taitensis</i>	Win mig	F coll. Oct 1926.	4,
10.	Common Golden Whistler	<i>Pachycephala pectoralis ornata</i>	Res bre	P, 8+ spec taken 1927	1,2,
11.	Cardinal Myzomela	<i>Myzomela cardinalis</i>	Res bre	P, Spec. AMNH	1,6,

### KEY

#### Ecological status

**End** = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

#### Abundance

**R** = rare, **S** = scarce, **UC** = uncommon, **MC** = moderately common, **LC** = locally common, **C** = common, **VC** = very common, **A** = abundant **P** = present.

**Colour coding** applies only to one species on a particular list. eg. An abundance comment in **red** for the Brown Booby refers to the Ref # in **Red** for that species only.

**Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.**

### REFERENCES

- Mayr, E. 1945. **Birds of the South-west Pacific**. Wheldon Press. New York.
- Mayr, E. 1932. 27. Birds collected during the Whitney South Sea Expedition 21. Notes on Thickheads (Pachycephala) from Polynesia [sic]. *Am. Mus. Novit.* **522**: 1-22.
- Stickney, E. H. 1943. Birds collected during the Whitney South Sea Expedition. 53. Northern Shore birds in the Pacific. *Am. Mus. Novit.* **1248**: 1-9.
- Bogert, C. 1937. Birds Collected during the Whitney South Sea Expedition. 34. The distribution and migration of the Long-tailed Cuckoo (*Urodynamis taitensis* Sparrman). *Am. Mus. Novit.* **933**: 1-12.
- Mayr, E. & D. Amadon. 1941. Birds collected during the Whitney South Sea Expedition. 46. Geographical variation in *Demigretta sacra* (Gmelin). *Am. Mus. Novit.* **1144**: 1-11.
- Mayr, E. 1932. Birds collected during the Whitney South Sea Expedition. 18. Notes on Meliphagidae from Polynesia and the Solomon Islands. *Am. Mus. Novit.* **516**: 1-30.
- Amadon, D. 1943. Birds collected during the Whitney South Sea Expedition. 52. Notes on some non-passerine genera, 3. *Amer. Mus. Novit.* **1237**, 1-22.
- Bourne, W.R.P. in Dutson, G. 2001. New distributional ranges for melanesian birds. *Emu* 101, 237-248.
- Dutson, G. 2001. New distributional ranges for Melanesian birds. *Emu* **101**: 237-248.

