

Narega Is. Bird Checklist

Bismarck Sea, PNG.

04 32 48s 149 06 44e

Compiled by M.K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

(for contact, re-type e-mail address)

| # | Common Name | Scientific Name | Ecol. Status | Abundance | References |
|----|-----------------------|-------------------------------|--------------|---|------------|
| 1. | Streaked Shearwater | <i>Calonectris leucomelas</i> | Sum mig | 100 obs 81 km to NW 13.1.1986. | 1. |
| 2. | Heinroth's Shearwater | <i>Puffinus heinrothi</i> | Res? | 1 obs 81 km NNW 13.9.1990, 5 obs 80 km NNW 14.9.1990, | 1. |

KEY

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

R = rare, **S** = scarce, **UC** = uncommon, **MC** = moderately common, **LC** = locally common, **C** = common, **VC** = very common, **A** = abundant **P** = present.

Colour coding applies only to one species on a particular list. eg. An abundance comment in **red** for the Brown Booby refers to the Ref # in **Red** for that species only.

Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

REFERENCES

1. Cheshire, N. 2010. Procellariiformes observed around Papua New Guinea including the Bismarck Archipelago from 1985 to 2007. South Australian Ornithologist 36, 9-24.